

Differential calculus on a novel cross-product quantum algebra

Deepak Parashar

Department of Mathematics, University of Wales Swansea, Singleton Park, Swansea SA2 8PP, United Kingdom

E-mail: D.Parashar@swansea.ac.uk

URL: <http://www-maths.swan.ac.uk/staff/dp/>

Abstract. We investigate the algebro-geometric structure of a novel two-parameter quantum deformation which exhibits the nature of a semidirect or cross-product algebra built upon $GL(2) \otimes GL(1)$, and is related to several other known examples of quantum groups. Following the R-matrix framework, we construct the $L^\pm$ functionals and address the problem of duality for this quantum group. This naturally leads to the construction of a bicovariant differential calculus that depends only on one deformation parameter, respects the cross-product structure and has interesting applications. The corresponding Jordanian and hybrid deformation is also explored.

1. The quantum algebra $\mathcal{A}_{r,s}$

The biparametric q -deformation $\mathcal{A}_{r,s}$ is defined [1] to be the semidirect or cross-product $GL_r(2) \rtimes \mathbb{C}[f, f^{-1}]$ built on the vector space $GL_r(2) \otimes \mathbb{C}[f, f^{-1}]$ where $GL_r(2) = \mathbb{C}[a, b, c, d]$ modulo the relations

$$\begin{aligned} ab &= r^{-1}ba, & bd &= r^{-1}db \\ ac &= r^{-1}ca, & cd &= r^{-1}dc \\ bc &= cb, & [a, d] &= (r^{-1} - r)bc \end{aligned} \quad (1)$$

and $\mathbb{C}[f, f^{-1}]$ has the cross relations

$$\begin{aligned} af &= fa, & cf &= sfc \\ bf &= s^{-1}fb, & df &= fd \end{aligned} \quad (2)$$

$\mathcal{A}_{r,s}$ can also be interpreted as a skew Laurent polynomial ring $GL_r[f, f^{-1}; \sigma]$ where σ is the automorphism given by the action of element f on $GL_r(2)$. If we let $A = GL_r(2)$ and $H = \mathbb{C}[f, f^{-1}]$, then A is a left H -module algebra and the action of f is given by

$$f \triangleright a = a, \quad f \triangleright b = sb, \quad f \triangleright c = s^{-1}c, \quad f \triangleright d = d \quad (3)$$

2. The dual algebra $\mathcal{U}_{r,s}$

Knowing properties of cross-product algebras [2, 3], we already know that the algebra dual to $\mathcal{A}_{r,s}$ would be the cross-coproduct coalgebra $\mathcal{U}_{r,s} = U_r(gl(2)) \rtimes \mathbb{C}[[\phi]]$ with ϕ as an element dual to f . As a vector space, the dual is $\mathcal{U}_{r,s} = U_r(gl(2)) \otimes U(u(1))$. Now, the duality relation between $\langle GL_r(2), U_r(gl(2)) \rangle$ is already well-known [4], while that between $\langle \mathbb{C}[f, f^{-1}], U(u(1)) \rangle$ is given by $\langle f, \phi \rangle = 1$, i.e., $U(u(1)) = \mathbb{C}[[\phi]]$. More precisely, we work

algebraically with $\mathbb{C}[s^\phi, s^{-\phi}]$ where $\langle f, s^\phi \rangle = s$. This induces duality on the vector space tensor products, the left action dualises to the left coaction, and this results in the dual algebra being a cross-coproduct $\mathcal{U}_{r,s} = U_r(gl(2)) \times_s \mathbb{C}[[\phi]]$. Let us recall that $U_r(gl(2))$, the algebra dual to $GL_r(2)$, is isomorphic to the tensor product $U_r(sl(2)) \otimes \tilde{U}(u(1))$ where $U_r(sl(2))$ has the usual generators $\{H, X_\pm\}$ and $\tilde{U}(u(1)) = \mathbb{C}[[\xi]] = \mathbb{C}[r^\xi, r^{-\xi}]$ with ξ central. Therefore, $\mathcal{U}_{r,s}$ is nothing but $U_r(sl(2))$ and two central generators ξ and ϕ , where ξ is the generating element of $\tilde{U}(u(1))$ and ϕ is the generating element of $U(u(1))$. Also note that s^ϕ is dually paired with the element f of $\mathcal{A}_{r,s}$.

3. R -matrix relations

In the quantum group language, $\mathcal{A}_{r,s}$ is understood as a novel Hopf algebra [5, 6] generated by $\{a, b, c, d, f\}$ arranged in the matrix form

$$T = \begin{pmatrix} f & 0 & 0 \\ 0 & a & b \\ 0 & c & d \end{pmatrix} \quad (4)$$

with the labelling 0, 1, 2, and $\{r, s\}$ are the two deformation parameters. The R -matrix

$$R = \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & S^{-1} & 0 & 0 \\ 0 & \Lambda & S & 0 \\ 0 & 0 & 0 & R_r \end{pmatrix} \quad (5)$$

is in block form, i.e., in the order (00), (01), (02), (10), (20), (11), (12), (21), (22) (which is chosen in conjunction with the block form of the T -matrix) where

$$R_r = \begin{pmatrix} r & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & \lambda & 1 & 0 \\ 0 & 0 & 0 & r \end{pmatrix}; \quad S = \begin{pmatrix} s & 0 \\ 0 & 1 \end{pmatrix}; \quad \Lambda = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}; \quad \lambda = r - r^{-1}$$

The RTT relations $RT_1T_2 = T_2T_1R$ (where $T_1 = T \otimes 1$ and $T_2 = 1 \otimes T$) then yield the commutation relations (1) and (2) between the generators. The Hopf algebra structure underlying $\mathcal{A}_{r,s}$ is $\Delta(T) = T \otimes T$, $\varepsilon(T) = 1$. The Casimir operator $\delta = ad - r^{-1}bc$ is invertible and the antipode is

$$S(f) = f^{-1}, \quad S(a) = \delta^{-1}d, \quad S(b) = -\delta^{-1}rb, \quad S(c) = -\delta^{-1}r^{-1}c, \quad S(d) = \delta^{-1}a \quad (6)$$

The quantum determinant $\mathcal{D} = \delta f$ is group-like but not central. There are several interesting features of this deformation [5, 6, 1] which cannot be mentioned here. In the R -matrix formulation of matrix quantum groups, a basic step is to construct functionals (matrices) L^+ and L^- which are dual to the matrix of generators (4) in the fundamental representation. These functions are defined by their value on the matrix of generators T

$$\langle (L^\pm)_b^a, T_d^c \rangle = (R^\pm)_{bd}^{ac} \quad (7)$$

where

$$(R^+)_{bd}^{ac} = c^+(R)_{db}^{ca} \quad (8)$$

$$(R^-)_{bd}^{ac} = c^-(R^{-1})_{db}^{ca} \quad (9)$$

and c^+ , c^- are free parameters. For $\mathcal{A}_{r,s}$ we make the following ansatz for the $L^\pm$ matrices:

$$L^+ = \begin{pmatrix} J & 0 & 0 \\ 0 & M & P \\ 0 & 0 & N \end{pmatrix} \quad \text{and} \quad L^- = \begin{pmatrix} J' & 0 & 0 \\ 0 & M' & 0 \\ 0 & Q & N' \end{pmatrix} \quad (10)$$

where

$$\begin{aligned} J &= s^{-\frac{1}{2}(F-A+D-1)} r^{\frac{1}{2}(F-A-D+1)}, & J' &= s^{-\frac{1}{2}(F-A+D-1)} r^{-\frac{1}{2}(F-A-D+1)} \\ M &= s^{-\frac{1}{2}(F-A-D+1)} r^{\frac{1}{2}(-F+A-D+1)}, & M' &= s^{-\frac{1}{2}(F-A-D+1)} r^{-\frac{1}{2}(-F+A-D+1)} \\ N &= s^{-\frac{1}{2}(F+A+D-1)} r^{\frac{1}{2}(-F-A+D+1)}, & N' &= s^{-\frac{1}{2}(F+A+D-1)} r^{-\frac{1}{2}(-F-A+D+1)} \\ P &= \lambda C, & Q &= -\lambda B \end{aligned}$$

and $\{A, B, C, D, F\}$ is the set of generating elements of the dual algebra $\mathcal{U}_{r,s}$. This is consistent with the action on the generators of $\mathcal{A}_{r,s}$ and gives the correct duality pairings. The commutation algebra is given by the RLL relations $R_{12}L_2^\pm L_1^\pm = L_1^\pm L_2^\pm R_{12}$, $R_{12}L_2^+ L_1^- = L_1^- L_2^+ R_{12}$, where $L_1^\pm = L^\pm \otimes \mathbf{1}$, $L_2^\pm = \mathbf{1} \otimes L^\pm$, and R_{12} is the same as (5). Finally, we obtain a single-parameter deformation of $U(gl(2)) \otimes U(u(1))$ as an algebra. Including the coproduct, we again obtain $\mathcal{U}_{r,s}$ as a semidirect product $U_r(gl(2)) \rtimes_s U(u(1))$.

4. Differential calculus on $\mathcal{A}_{r,s}$

The R -matrix procedure [7] is known to provide a natural framework to construct differential calculus on matrix quantum groups. We note here that the $\mathcal{A}_{r,s}$ deformation is not a full matrix quantum group, but an appropriate quotient of one (of multiparameter q -deformed $GL(3)$, to be precise). Nevertheless, it turns out that the constructive differential calculus methods [8] work equally well for such quotients. The bimodule Γ (space of quantum one-forms ω) is characterised by the commutation relations between ω and $a \in \mathcal{A} (\equiv \mathcal{A}_{r,s})$

$$\omega a = (\mathbf{1} \otimes g) \Delta(a) \omega \quad (11)$$

and the linear functional $g \in \mathcal{A}' (= \text{Hom}(\mathcal{A}, \mathbb{C}))$ is defined in terms of the $L^\pm$ matrices

$$g = S(L^+) L^- \quad (12)$$

Thus, in terms of components we have

$$\omega_{ij} a = [(\mathbf{1} \otimes S(l_{ki}^+ l_{jl}^-) \Delta(a)) \omega_{kl}] \quad (13)$$

using $L^\pm = l_{ij}^\pm$ and $\omega = \omega_{ij}$ where $i, j = 1..3$. From these relations, one can obtain the commutation relations of all the left-invariant one-forms with the generating elements of $\mathcal{A}$. The left-invariant vector fields χ_{ij} on $\mathcal{A}$ are given by the expression

$$\chi_{ij} = S(l_{ik}^+) l_{kj}^- - \delta_{ij} \varepsilon \quad (14)$$

The vector fields act on the generating elements as

$$\chi_{ij} a = (S(l_{ik}^+) l_{kj}^- - \delta_{ij} \varepsilon) a \quad (15)$$

Furthermore, using the formula $\mathbf{d}a = \sum_i (\chi_i * a) \omega^i$, we obtain the action of the exterior derivatives ($\mathbf{d} : \mathcal{A} \longrightarrow \Gamma$):

$$\mathbf{d}a = (r^{-2} - 1)a\omega^1 - \lambda b\omega^+ \quad (16)$$

$$\mathbf{d}b = \lambda^2 b\omega^1 - \lambda a\omega^- + (r^{-2} - 1)b\omega^2 \quad (17)$$

$$\mathbf{d}c = (r^{-2} - 1)c\omega^1 - \lambda d\omega^+ \quad (18)$$

$$\mathbf{d}d = \lambda^2 d\omega^1 - \lambda c\omega^- + (r^{-2} - 1)d\omega^2 \quad (19)$$

$$\mathbf{d}f = (r^{-2} - 1)f\omega^0 \quad (20)$$

where $\omega^0 = \omega_{11}, \omega^1 = \omega_{22}, \omega^+ = \omega_{23}, \omega^- = \omega_{32}, \omega^2 = \omega_{33}$. $\mathbf{d}\mathcal{A}$ generates Γ as a left $\mathcal{A}$ -module, and this defines a first-order differential calculus $(\Gamma, \mathbf{d})$ on $\mathcal{A}_{r,s}$. The calculus is bicovariant due to the coexistence of the left $(\Delta_L : \Gamma \longrightarrow \mathcal{A} \otimes \Gamma)$ and the right $(\Delta_R : \Gamma \longrightarrow \Gamma \otimes \mathcal{A})$ actions. Curiously, using the Leibniz rule it can be checked that

$$\mathbf{d}(af - fa) = 0, \quad \mathbf{d}(cf - sc) = 0, \quad \mathbf{d}(bf - s^{-1}fb) = 0, \quad \mathbf{d}(df - fd) = 0, \quad (21)$$

which is consistent with cross relations (2), and so the differential calculus also respects the cross-product structure of $\mathcal{A}_{r,s}$.

5. The Jordanian deformation $\mathcal{A}_{m,k}$

$\mathcal{A}_{r,s}$ can be contracted [6] (by means of singular limit of similarity transformations) to obtain a nonstandard or Jordanian analogue, say $\mathcal{A}_{m,k}$, with deformation parameters $\{m, k\}$ and the associated R -matrix is triangular. In analogy with $\mathcal{A}_{r,s}$, $\mathcal{A}_{m,k}$ can also be considered as the semidirect or cross-product $GL_m(2) \rtimes_k \mathbb{C}[f, f^{-1}]$ built upon the vector space $GL_m(2) \otimes \mathbb{C}[f, f^{-1}]$, where $GL_m(2)$ is itself a Jordanian deformation of $GL(2)$. Thus, $\mathcal{A}_{m,k}$ can also be interpreted as a skew Laurent polynomial ring $GL_m[f, f^{-1}; \sigma]$ where σ is the automorphism given by the action of element f on $GL_m(2)$.

6. Conclusions

The $\mathcal{A}_{r,s}$ and $\mathcal{A}_{m,k}$ deformations provide interesting new examples of cross-product quantum algebras, both of which have $GL(2) \otimes GL(1)$ as their classical limits. The differential calculus on $\mathcal{A}_{r,s}$ also has an inherent cross-product structure, embeds the calculus on $GL_q(2)$ and is also related to the calculus on $GL_{p,q}(2)$. It would be interesting to investigate the calculus on the Jordanian $\mathcal{A}_{m,k}$, and on the *hybrid/intermediate* [9] deformation obtained during the course of the contraction of $\mathcal{A}_{r,s}$ to $\mathcal{A}_{m,k}$.

References

- [1] Parashar D 2001 J. Math. Phys. 42 5431-5443
- [2] Majid S 1995 Foundations of Quantum Group Theory (Cambridge: CUP)
- [3] Klimyk A and Schmüdgen K 1997 Quantum Groups and Their Representations (Springer)
- [4] Sudbery A 1990 Proc. Workshop on Quantum Groups, Argonne (edited by Curtright T, Fairlie D and Zachos C) pp. 33-51
- [5] Basu-Mallick B 1994 hep-th/9402142
- [6] Parashar D and McDermott R J 2000 J. Math. Phys. 41 2403-2416
- [7] Faddeev L D, Reshetikhin N Y and Takhtajan L A 1990 Len. Math. J. 1 193-225
- [8] Jurčo B 1991 Lett. Math. Phys. 22 177-186; 1994 preprint CERN-TH 9417/94
- [9] Ballesteros A, Herranz F J and Parashar P 1999 J. Phys. A: Math. Gen. 32, 2369-2385